

Phytochemical profile, antioxidant activity and cytotoxicity against keratinocytes, fibroblasts and endothelial cells of picoplanktonic marine cyanobacteria

JANAÍNA MORONE^{1,2}, GRACILIANA LOPES^{1,2}, MARCO PRETO², VÍTOR VASCONCELOS^{1,2}, ROSÁRIO MARTINS^{2,3}

¹Faculty of Sciences, University of Porto, Porto, Portugal

²Interdisciplinary Centre of Marine and Environmental Research (CIIMAR/CIMAR), University of Porto, Matosinhos, Portugal

³Health and Environment Research Centre, School of Health, Polytechnic Institute of Porto, Porto, Portugal

Combining the increase demand for natural products in skin care formulations, and the bioactive arsenal of cyanobacteria, we aimed with this study to evaluate the potential of a 70% ethanolic extract of picocyanobacteria strains of the genera *Cyanobium* and *Synechocystis* for skin care applications. The cyanobacteria extract was analyzed for the phytochemical profile including carotenoid and phenolic content, for the antioxidant potential, and for the *in vitro* cytotoxicity against keratinocytes (HaCat), fibroblasts (3T3L1) and endothelial cells (hCMEC/D3). The total carotenoid content ranged from 162.43 to 383.89 $\mu\text{g g}^{-1}$ of dry biomass and the total phenolic content (TPC) from 1.09 to 2.45 mg GAE g^{-1} . Identified carotenoids consisted in zeaxanthin, lutein, canthaxanthin, echinenone and β -carotene, being zeaxanthin and lutein the most representative (49.82 and 79.08 $\mu\text{g g}^{-1}$, respectively). The antioxidant potential assessed by the DPPH \cdot and superoxide anion radical ($\text{O}_2\cdot^-$) scavenging assays resulted in an IC_{50} of 863.82 $\mu\text{g mL}^{-1}$ and 1275.86 $\mu\text{g mL}^{-1}$ respectively. An increase in cell viability was registered, particularly for fibroblasts and keratinocytes. From all the strains, both the species *Synechocystis salina* LEGE06099 and *Synechocystis salina* LEGE06155 evidenced an interesting potential for further exploitation.

Keywords: cyanobacteria, phytochemicals, antioxidant potential, cytotoxicity

Acknowledgments:

This work was done in the framework of the project BLUEHUMAN - BLUE biotechnology as a road for innovation on HUMAN's health aiming smart growth in Atlantic Area - EAPA_151/2016 of the Interreg Atlantic Area Programme funded by the European Regional Development Fund and ALGAVALOR - MicroALGAs: integrated production and valorization of biomass and its various applications, supported by the PORTUGAL 2020 through the European Regional Development Fund (ERDF).

This work was supported by the FCT Project UID/Multi/04423/2019 and by the Atlantic Interreg Project - EnhanceMicroAlgae - High added-value industrial opportunities for microalgae in the Atlantic Area (EAPA_338/2016).